



Fig. 5. Illustration for the proof of Theorem 3: Highlighting the basic topology of the agent paths that define the AGV-RAS employed in the relevant reduction from 3-SAT.

the context of some multi-vehicle traffic systems that are encountered in modern technological applications. In the process of deriving these results, we also developed a new proof for the NP-completeness of state safety in the class of L-SU-RAS, which is more concise and more lucid than the currently existing one. Two additional, similar problems that remain open, concern the computational complexity of state safety in (a) *dynamically routed* AGV and free-range vehicular systems, where a vehicle can be routed dynamically to any zone/cells neighboring its current location, and (b) in *closed* AGV and free-range vehicular systems where agents never retire from the domain of their motion, but they remain in it between the assignment of two consecutive trips. Furthermore, it is also interesting to identify any special structure under which the FREE-RANGE-RAS and AGV-RAS state safety problems, considered in this work, acquire polynomial complexity. Such a line of research can leverage and expand results of similar type for the L-SU-RAS state safety problem reported in [7, Ch.3].

REFERENCES

- [1] M. P. Groover, *Fundamentals of Modern Manufacturing: Materials, Processes and Systems*. Englewood Cliffs, NJ: Prentice Hall, 1996.
- [2] J. E. Hopcroft and J. D. Ullman, *Introduction to Automata Theory, Languages and Computation*. Reading, MA: Addison-Wesley, 1979.
- [3] A. Kobetski, "Optimal Coordination of Flexible Manufacturing Systems With Automatic Generation of Collision and Deadlock-Free Working Schedules," Ph.D. dissertation, Chalmers Univ. Technol., Goteborg, Sweden, 2008.
- [4] S. M. La Valle and S. A. Hutchinson, "Optimal motion planning for multiple robots having independent goals," *IEEE Trans. Robot. Autom.*, vol. 14, no. 6, pp. 912–925, Dec. 1998.
- [5] M. A. Lawley and S. A. Reveliotis, "Deadlock avoidance for sequential resource allocation systems: Hard and easy cases," *Int. J. FMS*, vol. 13, pp. 385–404, 2001.
- [6] S. A. Reveliotis, "Conflict resolution in AGV systems," *IIE Trans.*, vol. 32, no. 7, pp. 647–659, 2000.
- [7] S. A. Reveliotis, *Real-Time Management of Resource Allocation Systems: A Discrete Event Systems Approach*. New York: Springer, 2005.
- [8] S. A. Reveliotis and E. Roszkowska, "Conflict resolution in multi-vehicle systems: A resource allocation paradigm," in *Proc. IEEE CASE'08*, 2008, pp. 115–121.
- [9] E. Roszkowska, "High-level motion control for workspace sharing mobile robots," in *Robot Motion and Control*. New York: Springer, 2007, vol. 360, LNCIS, pp. 427–35.
- [10] E. Roszkowska and S. Reveliotis, "On the liveness of guidepath-based, zoned-controlled, dynamically routed, closed traffic systems," *IEEE Trans. Autom. Control*, vol. 53, no. 7, pp. 1689–1695, Jun. 2008.
- [11] N. Wu and M. Zhou, "Resource-oriented Petri nets in deadlock avoidance of AGV systems," in *Proc. ICRA'01*, 2001, pp. 64–69.

On Conditions for Convergence to Consensus

Jan Lorenz and Dirk A. Lorenz

Abstract—A new theorem on conditions for convergence to consensus of a multiagent time-dependent time-discrete dynamical system is presented. The theorem is build up on the notion of averaging maps. We compare this theorem to results by Moreau [6] (IEEE TRANSACTIONS ON AUTOMATIC CONTROL, vol. 50, no. 2, 2005) about set-valued Lyapunov theory and convergence under switching communication topologies. We give examples that point out differences of approaches including examples where Moreau's theorem is not applicable but ours is. Further on, we give examples that demonstrate that the theory of convergence to consensus is still not complete.

Index Terms—Averaging map, consensus protocol, multiagent systems, set-valued Lyapunov theory.

I. INTRODUCTION

In this technical note we analyze discrete dynamical systems of consensus formation as presented in the context of distributed computing [1], [2], flocking (e.g., of unmanned aerial vehicles) [3]–[5] and general as multi-agent coordination problems [6]–[8] (to mention just a few). The dynamical system may also be called 'agreement algorithm' or 'consensus protocol'. The convergence theorems of Moreau [6] together with the extensions of Angeli and Bliman [9] are the most general ones. The main theorem of Moreau states conditions for convergence to consensus under switching communication topologies. Convergence to consensus is there implied by 'global asymptotic stability of the set of equilibrium solutions with consensus as equilibrium points'. Conditions are on the one hand on the communication topologies in their time-evolution and on the other hand on the updating maps. Moreau applied a set-valued Lyapunov theory, which uses a set-valued function on the state space which is contractive with respect to the updating map. This implies convergence of the set to a singleton.

We contribute a similar but new approach based on the notion of an averaging map. Moreau deals with communication topologies by defining conditions on how many successive communication topologies must be regarded until the composition of these updating maps fulfills the contraction properties used to apply the set-valued Lyapunov theory. We skip the issue on changing communication topologies and deal directly with maps which fulfill a contraction property which is different from Moreau's.

Our theorem generalizes a result of Krause [10] by allowing arbitrary switching between different averaging maps but follow the same line of compactness, continuity and convexity arguments.

Section II presents the convergence result and possible extensions. Section III discusses the relations to two of Moreau's theorems in more detail. Section IV gives examples and counterexamples to show existing gaps in the theory of consensus algorithms. All proofs of lemmas and theorems are collected in Appendix A.

Manuscript received November 24, 2008; revised July 21, 2009. First published March 22, 2010; current version published July 08, 2010. Recommended by Associate Editor Z. Qu.

J. Lorenz is with the Chair of Systems Design, ETH Zurich, Zurich 8032, Switzerland (e-mail: post@janlo.de; URL: <http://www.janlo.de>).

D.A. Lorenz is with the Institute for Analysis and Algebra, Carl-Friedrich Gauß Department, TU Braunschweig, Braunschweig 38092, Germany (e-mail: d.lorenz@tu-braunschweig.de).

Digital Object Identifier 10.1109/TAC.2010.2046086

II. CONVERGENCE RESULT

We consider a dynamical system of the form

$$x(t+1) = f_t(x(t)) \quad (1)$$

with discrete time $t \in \mathbb{N}$. Dynamics take place in a $d \times n$ -dimensional space: We consider a set of agents $\underline{n} = \{1, \dots, n\}$ where each of them has coordinates in a d -dimensional set $S \subset \mathbb{R}^d$. Hence, the solutions of (1) have the form $x : \mathbb{N} \rightarrow S^n \subset \mathbb{R}^{d \times n}$. The individual coordinates of agent i at time $t \in \mathbb{N}$ is labeled $x^i(t) \in S$, and $x(t) \in S^n$ is called the *profile* at time $t \in \mathbb{N}$. Finally, the mappings f_t which govern the dynamics are of the form $f_t : S^n \rightarrow S^n$. We denote the component functions by f_t^i .

To state our main result on convergence of such systems to consensus we introduce the following notations. An element $x \in S^n$ is called *consensus* if all d -dimensional coordinates x^i have the same value, i.e., there exists a vector $\gamma \in S$ such that $x^i = \gamma$ for $i \in \underline{n}$. By $\text{conv}_{i \in \underline{n}} x^i$ we define the convex hull of the vectors x^1, \dots, x^n .

The core notion in this note is an 'averaging map'. We build the definition of an averaging map on a generalized convex hull. Consider a continuous function $y : S^n \rightarrow S^m$ which maps a profile to a certain set of m vectors $y(x) = (y^1(x), \dots, y^m(x))$ such that for all $x \in S^n$ and all $i \in \underline{n}$ it holds $x^i \in \text{conv}_{j \in \underline{m}} y^j(x)$. We call such a function y a *generalized barycentric coordinate map* and we call $\text{conv}_{j \in \underline{m}} y^j(x)$ the *y -convex hull* of the vectors x^1, \dots, x^n . (We call y 'generalized' because it needs not be a bijective transformation.) So, a y -convex hull is a set-valued function from S^n to the compact and convex subsets of S . We call a set y -convex, if it is the union of the y -convex hulls of all n of its points. Examples for y -convex hulls include the convex hull itself, and the multidimensional interval $[\min_{i \in \underline{n}} x^i, \max_{i \in \underline{n}} x^i]$ (with \min and \max applied componentwise). For the first it holds $m = n$ for the second $m = 2^d$. Many other examples fit into this setting: the smallest interval for any basis of \mathbb{R}^d [9, Example 2], or smallest polytope with faces parallel to a set of $k \geq d+1$ hyperplanes [9, Example 3] containing x^1, \dots, x^n (the generalized barycentric coordinates are then the extreme points of the polytope, perhaps with multiples to have a constant m). Now, we define the central notion of this technical note.

Definition 2.1: Let $S \subset \mathbb{R}^d$, $y : S^n \rightarrow S^m$ be a generalized barycentric coordinate map such that S is y -convex. A mapping $f : S^n \rightarrow S^n$ is called a *y -averaging map*, if for every $x \in S^n$ it holds

$$\text{conv}_{i \in \underline{m}} y^i(f(x)) \subset \text{conv}_{i \in \underline{m}} y^i(x). \quad (2)$$

Furthermore, a *proper y -averaging map* is a y -averaging map, such that for every $x \in S^n$ which is not a consensus, the above inclusion is strict.

A y -averaging map maps a profile x into its y -convex hull. Furthermore, the y -convex hull of the new profile $f(x)$ lies in the y -convex hull of the vectors x^1, \dots, x^n . Hence, we may also work with the y -convex hull of the initial profile $x(0)$ instead of the set S . Sometimes it is useful to look at the contraposition of the definition of proper: If equality holds in (2) this implies that x is a consensus. In the following we may omit 'y' when we mention an averaging map, but for an averaging map the definition of y is a prerequisite. The best proxy for the mind is $y = \text{id}$.

Since we are going to consider families of averaging maps we introduce the concept of equiproper averaging maps. To this end, we need the Hausdorff distance on the set of compact subsets of a metric space (X, d) . The distance of a point $x \in X$ and a nonempty compact set $C \subset X$ is defined as $d(x, C) := \min_{c \in C} d(x, c)$. Let $B, C \subset X$ be nonempty and compact, then the *Hausdorff distance* is defined as

$$d_H(B, C) := \max \left\{ \max_{b \in B} d(b, C), \max_{c \in C} d(c, B) \right\}.$$

Equivalently, one can say that the Hausdorff distance is the smallest ε such that the ε -neighborhood of B contains C and the ε -neighborhood of C contains B . It is easy to see that $d_H(B, C) = 0$ holds if and only if $B = C$. In the special case $B \subset C \subset S \subset \mathbb{R}^d$ it holds

$$d_H(B, C) = \max_{b \in B} d(b, C) = \max_{b \in B} \min_{c \in C} \|b - c\|. \quad (3)$$

Definition 2.2: Let y be a generalized barycentric coordinate map and let F be a family of proper y -averaging maps. F is called *equiproper*, if for every $x \in S^n$ which is not a consensus, there is $\delta(x) > 0$ such that for all $f \in F$

$$d_H \left(\text{conv}_{i \in \underline{m}} y^i(f(x)), \text{conv}_{i \in \underline{m}} y^i(x) \right) > \delta(x). \quad (4)$$

Now we state a lemma which says that the family of equiproper y -averaging maps is closed under pointwise limits.

Lemma 2.3: Let f_t be a sequence of y -averaging maps forming an equiproper family of y -averaging maps such that $f_t \rightarrow g$ pointwise. Then g is a proper y -averaging map.

Now we are able to state our main theorem.

Theorem 2.4: Let $S \subset \mathbb{R}^d$, y be a generalized barycentric coordinate map such that S is y -convex, and F be an equicontinuous family of equiproper y -averaging maps on S^n . Then it holds for any sequence $(f_t)_{t \in \mathbb{N}}$ with $f_t \in F$ and any $x(0) \in S^n$ that the solution of (1) converges to a consensus, i.e., there exists $\gamma \in S$ such that for all $i \in \underline{n}$ it holds $\lim_{t \rightarrow \infty} x^i(t) = \gamma$.

Notice that the limit γ depends not only on the initial value $x(0)$ but also on the realization of the sequence $(f_t)_{t \in \mathbb{N}}$, however, γ depends continuously on the initial value if the sequence (f_t) is fixed as the following lemma and corollary show.

Lemma 2.5: Let (X, d) be a metric space and $f_t : X \rightarrow X$ be such that the solution of $x(t+1) = f_t(x(t))$ converge to some limit for every initial value $x(0) \in X$. Then the limit depends continuously on the initial value if $\{f_t\}$ is an equicontinuous family.

The following corollary is a direct consequence.

Corollary 2.6: Let the sequence (f_t) in the situation of Theorem 2.4 be fixed. Then the consensus value γ (which exists due to Theorem 2.4) depends continuously on the initial value.

Theorem 2.4 is a generalization of a theorem of Krause [10]. Krause's theorem is the special case when y is the identity and F contains only one proper averaging map. Notice that 'equi' in equiproper and equicontinuous can be omitted if F is a finite set. An easy extension is to allow F to contain also non-proper averaging maps (but at least one proper averaging map). Then the sequence $(f_t)_{t \in \mathbb{N}}$ has to contain a subsequence $(f_{t_s})_{s \in \mathbb{N}}$ of equiproper averaging maps to ensure convergence to consensus. This holds because then $\{g_s | g_s = f_{t_s} \circ \dots \circ f_{t_{s+1}}\}$ is an equiproper set of averaging maps for $s \in \mathbb{N}$. Notice that it is possible that a sequence of averaging maps contains a subsequence as above such that subcompositions g_s form an equiproper set, even when no f_t is proper. The easiest example is when F contains only one linear map which is determined by a row-stochastic square matrix which is regular but not scrambling (see Seneta [11]). For linear systems "row-stochastic" is equivalent to "being an averaging map" (with y the identity) and "scrambling" is equivalent to "proper." From the theory of nonnegative matrices we know that for each regular matrix there is an integer such that higher powers are scrambling.

In the spirit of [9] we state another generalization of Theorem 2.4 which deals with deformations of the hull. To this end, let $S, T \subset \mathbb{R}^d$ be compact and $\phi : T \rightarrow S$ be a homeomorphism. For a generalized barycentric coordinate map $y : S^n \rightarrow S^m$ we define the y, ϕ -hull as $\phi^{-1}(\text{conv}_{i \in \underline{m}} y^i(\phi(x)))$. Now, a y, ϕ -averaging map g is defined analogous to Definition 2.1

$$\phi^{-1} \left(\text{conv}_{i \in \underline{m}} y^i(\phi(g(x))) \right) \subset \phi^{-1} \left(\text{conv}_{i \in \underline{m}} y^i(\phi(x)) \right).$$

Note, that the y, ϕ -hull is not necessarily convex, see [9, Example 6]. The extension of the notions “proper” and “equiproper” is straightforward.

Theorem 2.7: Let $\phi : T \rightarrow S$ be continuous with Lipschitz continuous inverse and let y be a generalized barycentric coordinate map such that S is y -convex. Let G be a family of equicontinuous, equiproper y, ϕ -averaging maps on T^n . Then it holds for any sequence $(g_t)_{t \in \mathbb{N}}$ with $g_t \in G$ and any $x(0) \in T^n$ that the solution of $x(t+1) = g_t(x(t))$ converges to a consensus.

III. COMPARISON WITH MOREAU’S SET-VALUED LYAPUNOV THEORY AND MAIN THEOREM

Theorem 2.4 has similarities to Moreau’s set-valued Lyapunov Theorem [6, Theorem 4]. This theorem implies global asymptotic stability of the set of equilibrium solutions when there exists a set-valued function V on the state space, a measure for these sets μ , and a positive definite function β on the state space. Essentially it has to hold $V(f_t(x)) \subset V(x)$ and $\mu(V(f_t(x))) - \mu(V(x)) \leq -\beta(x)$. The best example to imagine is $V = \text{conv}$, and μ is the diameter of a set.

The set of equilibrium solutions for the dynamical system (1) under the conditions of Theorem 2.4 contains only all constant solutions on consensus vectors, due to the equiproperness of F . Given this set of equilibrium solutions, “global asymptotic stability of the set of equilibrium solutions” implies convergence to consensus. Convergence to consensus is thus a special case of the set-valued Lyapunov Theorem in [6]. To the best of our knowledge, it is the only case in which the theorem has been used so far.

Compared with our Theorem 2.4 the role of the set-valued map V is taken by the y -convex hull. So, we also deal with a general class of functions due to the various possible coordinate maps $y : S^n \rightarrow S^m$ —we only assume that m is finite. However, we do not need a general measure μ on these maps. The assumption $\mu(V(f_t(x))) - \mu(V(x)) < \beta(x)$ corresponds to $d_H(V(f_t(x)), V(x)) > \delta(x)$. This is a different condition and often weaker, as for example in the case where Moreau specifies it to proof his main Theorem [6, Theorem 2]. There μ is the diameter of $V(x)$ (which he specifies as the $\text{conv}(x)$).

Theorem 2.4 has also similarities to Moreau’s main theorem [6, Theorem 2]. This theorem is more specific than Theorem 2.4 by incorporating switching communication topologies. Its main drawback is that it relies very much on convex hulls (see [9] for a method to overcome this drawback). Our result generalizes to convex hulls of generalized coordinate maps. Further on, in Moreau’s theorem agents are forced to move into the relative interior of the convex hull (respecting the communication topology). Specifically, this implies that agents have to leave all extreme points of the convex hull (of agents in its neighborhood) after one iteration. Our theorem needs only agents at one arbitrary extreme point (of the global y -convex hull) to leave it towards the interior after one iteration. This is implied by properness of averaging maps. The assumption ‘equiproper’ in our theorem finds its analog in

Moreau’s theorem by assuming that the sets $e_k(\mathcal{A}(t))(x)$ are chosen independently of t .¹

Summarizing the above one can say that both Moreau’s theorem and Theorem 2.4 are similar. However, the assumptions as well as the methods of proof are different. On the one hand we do not incorporate switching communication topologies explicitly, but on the other hand we need weaker conditions for the updating maps f_t . Further on, we generalized to y -convex hulls and are also able to incorporate the extensions of Moreau’s theorem by Angeli and Bliman [9] to overcome the restriction to convex sets. Moreover, the notion of a (equi-)proper y -averaging map allows a systematic and structured treatment of consensus algorithms (see e.g., the results in Lemma 2.3 and Lemma 2.5). Hence, Theorem 2.4 together with 2.7 give an alternative approach to the analysis of consensus protocols whose applicability is illustrated by examples in the next section.

IV. EXAMPLES AND COUNTEREXAMPLES

In this section we present counterexamples (Examples 4.1–4.3) to point that the existing theory, including our Theorem 2.4, delivers no sharp results on convergence to consensus. We also give examples which show cases, where our theorem is applicable but Theorem 2 of Moreau is not (Examples 4.4–4.6).

Continuity, for instance, is not necessary for convergence to consensus since there are discontinuous proper averaging maps which converge to consensus (one may take different averaging maps on different subdomains of S). On the other hand discontinuity may destroy convergence to consensus even for proper averaging maps (see [12, Section 3.1] for examples for this phenomenon).

The next two examples illustrate the role of equiproperness.

Example 4.1 (Non-Equiproper not Leading to Consensus): Let

$$f_t(x^1, x^2) := \left(\left(1 - \frac{1}{4^t}\right)x^1 + \frac{1}{4^t}x^2, \frac{1}{4^t}x^1 + \left(1 - \frac{1}{4^t}\right)x^2 \right).$$

It is easy to see that for $t \geq 1$ and $x(1) = (0, 1)$ it holds that $x^1(t) < 1/3$ and $x^2(t) > 2/3$. Obviously, $\{f_t | t \in \mathbb{N}\}$ is not equiproper because f_t converges to the identity as $t \rightarrow \infty$.

Example 4.2 (Non-Equiproper Leading to Consensus): Let

$$f_t(x^1, x^2) := \left(\left(1 - \frac{1}{t}\right)x^1 + \frac{1}{t}x^2, x^2 \right).$$

This example is not equiproper, because f_t converges to the identity for $t \rightarrow \infty$. Thus, Theorem 2.4 does not apply, but for $t \geq 2$ and any $x(2) \in (\mathbb{R})^2$ the system $x(t+1) = f_t(x(t))$ has the solution $x(t) = (1/t - 1x^1(2) + t - 2/t - 1x^2(2), x^2(2))$ and thus converges to consensus at $x^2(2)$. Note that the convergence is not at an exponential rate. Convergence to consensus in the last example can also not be ensured by Moreau’s theorems.

The next example illustrates the role of equicontinuity and is inspired by bounded confidence [13].

1) Example 4.3 (Vanishing Confidence): Let $f_t : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with

$$f_t^i(x) := \frac{\sum_{j=1}^n D_t(|x^i - x^j|)x^j}{\sum_{j=1}^n D_t(|x^i - x^j|)}$$

¹Here the matrix $\mathcal{A}(t)$ is the arbitrarily chosen communication topology at time t and x is a given state. The set $e_k(\mathcal{A}(t))(x)$ is a subset of the relative interior of the convex hull of the neighbors of k (including k) in the current communication topology, and it determines the set where the state of node k has to remain in after one iteration. So, e_k has to be fixed for a given communication topology and a certain state regardless of the chosen updating map $f(t, \dots)$. This is in analogy to equiproper which implies the existence of a minimal Hausdorff distance $\delta(x)$ after one iteration for a given state but all possible averaging maps.

and $D_t : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$. Now, f_t is an averaging map for any choice of D_t . Further on, f_t is continuous if D_t is, and f_t is proper if D_t is strictly positive. We chose $D_t(y) := e^{-(y/\varepsilon)^t}$ as a sequence of functions which has the cutoff function as pointwise limit function. Hence, D_t is continuous but $\{D_t | t \in \mathbb{N}\}$ is not equicontinuous. For $x(0) = (0, 8)$, $\varepsilon = 1$ the process $x(t) = f_t(x(t))$ does not converge to consensus although only proper averaging maps are involved. Rough estimates show that $|x^1(t) - x^2(t)| \geq 4$.

For other settings convergence under vanishing confidence is possible, as numerical examples in [12] show.

The following examples are to show limitations of Moreau's Theorem 2 and how Theorem 2.4 can be applied to show convergence to consensus.

Example 4.4 (Rendezvous Problem With Watergun Sensors): We consider a version of the Rendezvous Problem [14] where n agents are to locate themselves decentralized at the same position in twodimensional space. Each agent has three waterguns, an *activation gun* and two *search guns*. Agents can perceive from which kind of gun they were hit and can respond (e.g., acoustically). The search gun is used as a sensor to check if there is at least one other agent in direction $\alpha \in [0, 2\pi[$. The activation gun is used to activate other agents. When another agent responds to a shot by the activation gun, the shooting agent switches to standby (only responding if hit). With two search guns an agent can particularly perform a move into direction $\beta \in [0, 2\pi[$ under

Rule (*): Move until either the position of an other agent is reached or until there is an agent in the directions $\beta + \pi/2$ or $\beta - \pi/2$. (Move while constantly shooting left and right with search gun until someone is hit.)

Initially the n agents are located at different positions in space and the multi-agent protocol is started from the outside by activating one agent. Whenever an agent is activated it executes the following program:

search gun all around shot, detect A as set of all directions where agents are

select α, γ such that for all $-1 \leq c \leq 1$ it holds $(\alpha + c\gamma) \bmod 2\pi \notin A$ and γ maximal

if $\gamma \geq \pi/2 + \pi/n$ **then**

tie agents at same position to move together

move direction $\beta = \alpha + 2\pi \bmod 2\pi$ with rule (*)

end if

activation gun all around shot (random start) until someone hit

if no one hit **then**

give signal 'consensus found!'

end if

The protocol ensures that always only one agent is activated when an agent finishes its action unless consensus is found. It also always leads to the movement of an agent after some time unless consensus is found, because for every configuration there is always at least one agent whose position is an extreme point of the convex hull such that the exterior angle of the convex hull is larger than $\pi + 2\pi/n$ and thus $\gamma \geq \pi/2 + \pi/n$. This is because $\pi + 2\pi/n$ is the exterior angle of a regular polygon with n edges, which is the 'worst case'-polygon. It is 'worst case', because it has from all polygons with n edges the largest minimal exterior angle. Thus, the random search for an agent which finds a direction α always ends successfully unless consensus

is reached. So, the protocol leads to a series of actions which either continues forever including movements forever or finishing when consensus is reached. We group actions to form a series of updating maps f_t . We group by the following rule: Starting with the first action we collect actions in the same group until an agent is found which moves. The next updating map f_2 is formed analog starting with the next action, and so on. Thus we have a series of update maps.

It is simple to see that the series of updating maps f_1, f_2, \dots fulfills the conditions of Theorem 2.4 with y the identity. Every f_t is an averaging map because by definition the movement of agents goes into the convex hull or along its border and stops before the convex hull is left. It is equiproper, because for each x there are only as many possible updating maps as their convex hull has extreme points. Thus, there is a $\delta(x) > 0$ by taking the minimum over this finite set of possible updating maps. Every f_t is continuous in x when we regard all agents which have the same position as one agent. Equicontinuity at x again follows from finiteness of the possible updating maps.

Thus, the protocol in Example 4.4 leads to convergence to consensus. This can not be shown by applying Moreau's Theorem 2 because the movements cannot be easily encoded in terms of communication topologies. One could try to specify it in terms of communication topologies by stating that the moving agent has agents at the detected directions in A as its set of neighbors. But even then the conditions of Moreau's Assumption 1 (especially number 3) need not be fulfilled and a node connected to all other nodes across time intervals of length T need not exist as necessary for Moreau's Theorem 2.

Example 4.5 (Nonlinear Proper Averaging Map): Let

$$\begin{aligned} f_1(x) &= x_1, & f_2(x) &= a(l)x_1 + (1 - a(l))x_2, \\ f_3(x) &= \frac{1}{5}x_2 + \frac{4}{5}x_3 \end{aligned}$$

where $l = \text{dist}(x_3, \text{line passing through } x_1 \text{ and } x_2)$ and a is continuous and decreasing from $1/2$ to 0 in $[0, 1]$ and zero otherwise. In this example agent 3 moves towards agent 2 while agent 2 moves towards agent 1 only if agent 3 is close to a stripe around the line through agent 2 and agent 1.

Examples of this kind can be formulated in terms of communication topologies as Moreau's Theorem 2 needs them, but the existence of a uniform bound for the length of intercommunication intervals T is not easily at hand.

Example 4.6 (Non-Arithmetic Means): We define $g_1, g_2, g_3, g_4 : (\mathbb{R}^d)^3 \rightarrow \mathbb{R}^d$ by $g_1(x) := \max\{x^1, x^2, x^3\}$, $g_2(x) := 1/3(x^1 + x^2 + x^3)$, $g_3(x) := \sqrt[3]{x^1 x^2 x^3}$ and $g_4(x) := \min\{x^1, x^2, x^3\}$ with all computations componentwise. Further on let $f^{\sigma_1 \sigma_2 \sigma_3} : (\mathbb{R}^d)^3 \rightarrow (\mathbb{R}^d)^3$ with $f^{\sigma_1 \sigma_2 \sigma_3} := (g_{\sigma_1}, g_{\sigma_2}, g_{\sigma_3})$. It is easy to verify, that the family of all $f^{\sigma_1 \sigma_2 \sigma_3}$ where 1 and 4 are not both in $(\sigma_1, \sigma_2, \sigma_3) \in \{1, 2, 3, 4\}^3$ is an equicontinuous set of y -averaging maps, when the y -convex hull is the interval $[\min_{i \in \mathbb{N}} x^i, \max_{i \in \mathbb{N}} x^i]$. Equipropert is implied by finiteness. Thus convergence to consensus is ensured by Theorem 2.4. Moreau's theorem is not applicable because $f^{\sigma_1 \sigma_2 \sigma_3}$ is not a convex hull averaging map if some σ_i is 1, 3 or 4 (since the componentwise min or max and the geometric mean are in general not contained in the convex hull).

Krause [10] shows another example where Moreau's theorem does not imply convergence: Assume three agents in two dimensional space. In each iteration every agent takes the mean value of the two other agents. Hence, no agent moves into the relative interior of the convex hull but these maps are still proper averaging maps and Theorem 2.4 applies.

APPENDIX
 PROOFS

Proof of Lemma 2.3: First we show that g is an averaging map. Take $x \in S^n$ and let $\varepsilon > 0$. Due to the pointwise convergence of $(f_t)_i$ to g_i and uniform continuity of y there is t_0 such that for all $t > t_0$ it holds $\|y^i(f_t(x)) - y^i(g(x))\| < \varepsilon$. Due to $y^i(f_t(x)) \in \text{conv}_{i \in \underline{m}} y^i(x)$ it follows that the maximal distance of $y^i(g(x))$ to $\text{conv}_{i \in \underline{m}} y^i(x)$ is less than ε , and thus $y^i(g(x)) \in \text{conv}_{i \in \underline{m}} y^i(x)$ because $\text{conv}_{i \in \underline{m}} y^i(x)$ is closed.

We show that g is proper. To this end, let $x \in S^n$ be not a consensus. We have to show that there is $z^* \in \text{conv}_{i \in \underline{m}} y^i(x)$ but $z^* \notin \text{conv}_{i \in \underline{m}} y^i(g(x))$. (Note that $z^* \in S$, while $x \in S^n$ and $y(x) \in S^m$.) We know that there is for each $t \in \mathbb{N}$ an $z(t) \in \text{conv}_{i \in \underline{m}} y^i(x)$ with $z(t) \notin \text{conv}_{i \in \underline{m}} y^i(f_t(x))$. According to the equiproper property it can be chosen such that the distance of $z(t)$ to $\text{conv}_{i \in \underline{m}} y^i(f_t(x))$ is bigger than $\delta(x)/2 > 0$ for all $t \in \mathbb{N}$. Further on, we know that the set difference $\text{conv}_{i \in \underline{m}} y^i(f_t(x)) \setminus \text{conv}_{i \in \underline{m}} y^i(x)$ is non empty and bounded, thus there is a subsequence t_s such that $z(t_s)$ converges to a $z^* \in \text{conv}_{i \in \underline{m}} y^i(x)$. Because of the construction it also holds $z^* \notin \text{conv}_{i \in \underline{m}} y^i(g(x))$. ■

Proof of Theorem 2.4: The idea of the proof is the following: We define $C(t) := \text{conv}_{i \in \underline{m}} y^i(x(t))$ which is convex and compact. It holds $C(t+1) \subset C(t)$ because of the averaging property and $C := \bigcap_{t=0}^{\infty} C(t) \neq \emptyset$ because of compactness. In the following we will show that C is a singleton, and that for all $i \in \underline{n}$ the sequences $x^i(t)$ converge to it. This will be done in three main steps, but first we note that because of compactness of $C(0)^n$ there is a subsequence t_s and $c := (c^1, \dots, c^n) \in C(0)^n$ such that $\lim_{s \rightarrow \infty} x(t_s) = c$.

1) We show that $C = \text{conv}_{i \in \underline{m}} y^i(c)$. To accept “ \supset ” see that for all $t_s \geq t$ there is $x^i(t_s) \in C(t)$ and thus $c^i \in C(t)$. This implies $c^i \in C$ because all the $C(t)$ are closed.

To show “ \subset ” let $x \in C$ and $\varepsilon > 0$. Because of uniform continuity of y there is $\eta > 0$ such that for every $x' \in S$ with $\|c - x'\| < \eta$ it holds $\|y(c) - y(x')\| < \varepsilon$. Further on, there is s_0 such that for all $s \geq s_0$ it holds $\|x(t_s) - c\| < \eta$. This implies for every $i \in \underline{m}$ that $\|y^i(x(t_s)) - y^i(c)\| < \varepsilon$.

Obviously, $x \in C(t_{s_0})$. Thus, there exist convex coefficients $a_1, \dots, a_m \in \mathbb{R}_{\geq 0}^m$ such that $x = \sum_{i=1}^m a_i y^i(x(t_{s_0}))$. Now we can conclude

$$\begin{aligned} \left\| x - \sum_{i=1}^m a_i y^i(c) \right\| &= \left\| \sum_{i=1}^m a_i (y^i(x(t_{s_0})) - y^i(c)) \right\| \\ &\leq \sum_{i=1}^m \|y^i(x(t_{s_0})) - y^i(c)\| = m\varepsilon. \end{aligned}$$

It follows that $x \in \text{conv}_{i \in \underline{m}} y^i(c)$ because $\text{conv}_{i \in \underline{m}} y^i(c)$ is closed.

2) The next step is to show that c is a consensus, i.e., $c^1 = \dots = c^n$. The family F is uniformly equicontinuous and for all $x \in X$ it holds that $\{f(x) \mid f \in F\}$ is bounded (and thus relatively compact) because all the f are averaging maps. So, due to the theorem of Arzelà-Ascoli, F is relatively compact. Thus, there is a subsequence t_{s_r} such that $f_{t_{s_r}}$ converges uniformly to a continuous limit function g for $r \rightarrow \infty$. Due to Lemma 2.3 we also know that g is a proper averaging map. In two substeps we show that $\text{conv}_{i \in \underline{m}} y^i(g(c)) = \text{conv}_{i \in \underline{m}} y^i(c)$ which implies that c is a consensus:

a) We show that for all $i \in \underline{n}$ it holds $\lim_{r \rightarrow \infty} f_{t_{s_r}}(x_{t_{s_r}}) = g(c)$. We know that $f_{t_{s_r}} \rightarrow g$ uniformly and that $x^i(t_{s_r}) \rightarrow c$. Now we estimate

$$\begin{aligned} \|f_{t_{s_r}}(x(t_{s_r})) - g(c)\| &\leq \|f_{t_{s_r}}(x(t_{s_r})) - f_{t_{s_r}}(c)\| \\ &\quad + \|f_{t_{s_r}}(c) - g(c)\|. \end{aligned}$$

Both terms on the right hand side can be smaller than $\varepsilon/2$ for any ε for large enough r because of the continuity of $f_{t_{s_r}}$ and the uniform convergence $f_{t_{s_r}} \rightarrow g$.

b) We show $\text{conv}_{i \in \underline{m}} y^i(g(c)) = \text{conv}_{i \in \underline{m}} y^i(c)$. “ \subset ” holds because g is an averaging map (see 2a). To show “ \supset ” let $x \in \text{conv}_{i \in \underline{m}} y^i(c)$. Thus, for all r it holds $x \in C \subset C(t_{s_r}+1)$ and thus there exist convex coefficients with convex combination $x = \sum_{i=1}^m a_i(r) y^i(x(t_{s_r}+1))$. Now, $(a_1(r), \dots, a_m(r))_{r \in \mathbb{N}}$ is a sequence in the compact set of convex coefficients and thus there is a subsequence r_q such that $\lim_{q \rightarrow \infty} (a_1(r_q), \dots, a_m(r_q)) = (a_1^* \dots a_m^*)$. Now due to 2c and continuity of y it holds

$$\begin{aligned} x &= \sum_{i=1}^m \lim_{q \rightarrow \infty} a_i(r_q) \lim_{q \rightarrow \infty} y^i(x(t_{s_{r_q}}+1)) \\ &= \sum_{i=1}^m a_i^* y^i(g(c)). \end{aligned}$$

Thus, $x \in \text{conv}_{i \in \underline{m}} y^i(g(c))$.

This implies that c is a consensus, because g is a proper averaging map.

3) Finally, we show that for each $i \in \underline{n}$ the sequence $(x^i(t))_{t \in \mathbb{N}}$ (and not only subsequences) converges to $\gamma := c^1 = \dots = c^n$ for $t \rightarrow \infty$. We know that for $\varepsilon > 0$ there is a r_0 such that for each $i \in \underline{n}$ it holds $\|y^i(x(t_{s_{r_0}})) - \gamma\| < \varepsilon$. Further on, for $t \geq t_{s_{r_0}}$ it holds $x(t) \in C(t) \subset C(t_{s_{r_0}})$. Thus, for each $i \in \underline{n}$ there are convex combinations $x^i(t) = \sum_{j=1}^m a^j y^j(x(t_{s_{r_0}}))$. Now, we conclude for all $t > t_{s_{r_0}}$

$$\begin{aligned} \|x^i(t) - \gamma\| &= \left\| \sum_{j=1}^m a^j y^j(x(t_{s_{r_0}})) - \gamma \right\| \\ &\leq \sum_{j=1}^m \|x^j(t_{s_{r_0}}) - \gamma\| = m\varepsilon. \end{aligned}$$

This proves the theorem. ■

Proof of Lemma 2.5: Let $\varepsilon > 0$ and consider two initial values $x(0), \tilde{x}(0) \in S^n$ with corresponding limits $\gamma, \tilde{\gamma}$ respectively. We have to show that there exists $\delta > 0$ such that $d(x(0), \tilde{x}(0)) \leq \delta$ implies $d(\gamma, \tilde{\gamma}) \leq \varepsilon$.

We note that for every t it holds that

$$d(\gamma, \tilde{\gamma}) \leq d(\gamma, x(t)) + d(x(t), \tilde{x}(t)) + d(\tilde{\gamma}, \tilde{x}(t)).$$

We choose t_0 large enough, that

$$d(\gamma, x(t_0)) \leq \frac{\varepsilon}{3} \quad d(\tilde{\gamma}, \tilde{x}(t)) \leq \frac{\varepsilon}{3}.$$

Since $\{f_t\}$ is an equicontinuous family there exists $\eta > 0$ such that for every $t \in \mathbb{N}$ it holds that

$$d(x(t), \tilde{x}(t)) \leq \eta \Rightarrow d(f_t(x(t)), f_t(\tilde{x}(t))) \leq \varepsilon.$$

Since $x(t)$ and $\tilde{x}(t)$ solve $x(t+1) = f_t(x(t))$ we have recursively that for every t_0 there exists $\delta > 0$ such that

$$d(x(0), \tilde{x}(0)) \leq \delta \Rightarrow d(x(t_0), \tilde{x}(t_0)) \leq \frac{\varepsilon}{3}$$

which implies the claim. ■

Proof of Theorem 2.7: We define $f_t = \phi \circ g_t \circ \phi^{-1} : S^n \rightarrow S^n$. We show that $\{f_t \mid t \in \mathbb{N}\}$ is a family of equicontinuous, equiproper y -averaging maps on S^n . Equicontinuity and the fact that the f_t 's are

y -averaging maps are clear. To see equiproperness of f_t we note first that from equiproperness of g_t it follows:

$$\begin{aligned} d_H \left(\phi^{-1} \left(\underset{i \in \underline{m}}{\text{conv}} y^i (\phi(g_t(x))) \right), \phi^{-1} \left(\underset{i \in \underline{m}}{\text{conv}} y^i (\phi(x)) \right) \right) &\geq \delta(x) \\ \Rightarrow d_H \left(\phi^{-1} \left(\underset{i \in \underline{m}}{\text{conv}} y^i (f_t(\xi)) \right), \phi^{-1} \left(\underset{i \in \underline{m}}{\text{conv}} y^i (\xi) \right) \right) &\geq \delta(\phi^{-1}(\xi)) \end{aligned}$$

while the second line holds for all $\xi = \phi(x) \in S^n$ and $t \geq 0$. Due to (3) we can express the Hausdorff distance as

$$\begin{aligned} d_H \left(\phi^{-1} \left(\underset{i \in \underline{m}}{\text{conv}} y^i (f_t(\xi)) \right), \phi^{-1} \left(\underset{i \in \underline{m}}{\text{conv}} y^i (\xi) \right) \right) \\ = \max_{z \in \phi^{-1}(\underset{i \in \underline{m}}{\text{conv}} y^i (f_t(\xi)))} \min_{w \in \phi^{-1}(\underset{i \in \underline{m}}{\text{conv}} y^i (\xi))} \|z - w\| \\ = \max_{\phi(z) \in \underset{i \in \underline{m}}{\text{conv}} y^i (f_t(\xi))} \min_{\phi(w) \in \underset{i \in \underline{m}}{\text{conv}} y^i (\xi)} \|z - w\|. \end{aligned}$$

With this preparation we show equiproperness of the f_t 's

$$\begin{aligned} d_H \left(\underset{i \in \underline{m}}{\text{conv}} y^i (f_t(\xi)), \underset{i \in \underline{m}}{\text{conv}} y^i (\xi) \right) \\ = \max_{\zeta \in \underset{i \in \underline{m}}{\text{conv}} y^i (f_t(\xi))} \min_{\omega \in \underset{i \in \underline{m}}{\text{conv}} y^i (\xi)} \|\zeta - \omega\| \\ = \max_{\phi(z) \in \underset{i \in \underline{m}}{\text{conv}} y^i (f_t(\xi))} \min_{\phi(w) \in \underset{i \in \underline{m}}{\text{conv}} y^i (\xi)} \|\phi(z) - \phi(w)\| \\ \geq L \max_{\phi(z) \in \underset{i \in \underline{m}}{\text{conv}} y^i (f_t(\xi))} \min_{\phi(w) \in \underset{i \in \underline{m}}{\text{conv}} y^i (\xi)} \|z - w\| \\ \geq L\delta(\phi^{-1}(\xi)) \end{aligned}$$

where L is the Lipschitz constant of ϕ^{-1} . Now for $\xi(t) = \phi(x(t))$ it follows $\xi(t+1) = \phi(g_t(x(t))) = \phi(g_t(\phi^{-1}(\xi(t)))) = f_t(\xi(t))$. By virtue of Theorem 2.4, $\xi(t) \rightarrow c$ where $c \in S^n$ is a consensus and hence, $x(t) \rightarrow \phi^{-1}(c) \in T^n$ which is also a consensus. ■

REFERENCES

- [1] J. N. Tsitsiklis, "Problems in Decentralized Decision Making and Computation," Ph.D. dissertation, Dept. EECS, MIT, Cambridge, MA, 1984.
- [2] J. Tsitsiklis, D. Bertsekas, and M. Athans, "Distributed asynchronous deterministic and stochastic gradient optimization algorithms," *IEEE Trans. Autom. Control*, vol. AC-31, no. 9, pp. 803–812, Sep. 1986.
- [3] A. Jadbabaie, J. Lin, and A. Morse, "Coordination of groups of mobile autonomous agents using nearest neighbor rules," *IEEE Trans. Autom. Control*, vol. 48, no. 6, pp. 988–1001, Jun. 2003.
- [4] R. Saber and R. Murray, "Flocking with obstacle avoidance: Cooperation with limited communication in mobile networks," in *Proc. 42nd IEEE Conf. Decision Control*, Dec. 2003, vol. 2, pp. 2022–2028.
- [5] V. Blondel, J. Hendrickx, A. Olshevsky, and J. Tsitsiklis, "Convergence in multiagent coordination, consensus, and flocking," in *Proc. 44th IEEE Conf. Decision Control, & Eur. Control Conf. (CDC-ECC'05)*, Dec. 2005, pp. 2996–3000.
- [6] L. Moreau, "Stability of multiagent systems with time-dependent communication links," *IEEE Trans. Autom. Control*, vol. 50, no. 2, pp. 169–182, Feb. 2005.
- [7] W. Ren, R. Beard, and E. Atkins, "A survey of consensus problems in multi-agent coordination," in *Proc. Amer. Control Conf.*, Jun. 2005, vol. 3, pp. 1859–1864.
- [8] L. Xiao and S. Boyd, "Fast linear iterations for distributed averaging," *Syst. Control Lett.*, vol. 53, no. 1, pp. 65–78, 2004.
- [9] D. Angeli and P.-A. Bliman, "Stability of leaderless discrete-time multi-agent systems," *Math. Control, Signals, Syst. (MCSS)*, vol. 18, no. 4, pp. 293–322, Oct. 2006.
- [10] U. Krause, "Compromise, consensus, and the iteration of means," *Elemente der Mathematik*, vol. 64, no. 1, pp. 1–8, 2009.
- [11] E. Seneta, *Non-Negative Matrices and Markov Chains*. New York: Springer, 2006.
- [12] J. Lorenz, "Repeated Averaging and Bounded Confidence-Modeling, Analysis and Simulation of Continuous Opinion Dynamics" Ph.D. dissertation, Universität Bremen, Bremen, Germany, 2007.
- [13] R. Hegselmann and U. Krause, "Opinion dynamics and bounded confidence, models, analysis and simulation," *J. Artif. Societies Social Simul.*, vol. 5, no. 3, p. 2, 2002.
- [14] J. Lin, A. Morse, and B. Anderson, "The multi-agent rendezvous problem," in *Proc. 42nd IEEE Conf. Decision Control*, Dec. 2003, vol. 2, pp. 1508–1513.

Exponential Stabilization of a Class of Stochastic System With Markovian Jump Parameters and Mode-Dependent Mixed Time-Delays

Zidong Wang, Yurong Liu, and Xiaohui Liu

Abstract—In this technical note, the globally exponential stabilization problem is investigated for a general class of stochastic systems with both Markovian jumping parameters and mixed time-delays. The mixed mode-dependent time-delays consist of both discrete and distributed delays. We aim to design a memoryless state feedback controller such that the closed-loop system is stochastically exponentially stable in the mean square sense. First, by introducing a new Lyapunov-Krasovskii functional that accounts for the mode-dependent mixed delays, stochastic analysis is conducted in order to derive a criterion for the exponential stabilizability problem. Then, a variation of such a criterion is developed to facilitate the controller design by using the linear matrix inequality (LMI) approach. Finally, it is shown that the desired state feedback controller can be characterized explicitly in terms of the solution to a set of LMIs. Numerical simulation is carried out to demonstrate the effectiveness of the proposed methods.

Index Terms—Discrete time-delays, distributed time-delays, Markovian jumping parameters, mixed mode-dependent (MDD) time-delays, stochastic systems.

I. INTRODUCTION

It is now well known that time-delays are frequently encountered in practical systems such as engineering and biological systems, and their existence may induce instability, oscillation, and poor performances [1], [4], [5]. Time delays may also arise in several signal processing areas such as multipath propagation, telemanipulation systems, data communication in high-speed internet and network control systems [2]. According to the way time-delays occur, they can be classified as discrete (point) delays [16] and distributed delays [10]. In the past few years, considerable attention has been devoted to the robust stabilization and H_∞ control problem for linear and nonlinear time-delay systems, and a great number of papers have appeared on this general topic, see [2] for a survey.

Markovian jump systems (MJSs) involve both time-evolving and event-driven mechanisms, which can be employed to model the abrupt phenomena such as random failures and repairs of the components, changes in the interconnections of subsystems, sudden environment changes, etc. The issues of stability, stabilization, control and filtering have been well investigated, see e.g. [1], [3], [7], [9], [13], [15], [16], [19]. On another research forefront, since stochastic phenomenon typically exhibits in many branches of science and engineering applica-

Manuscript received July 17, 2009; revised November 08, 2009, February 09, 2010, March 06, 2010, and March 10, 2010. First published March 22, 2010; current version published July 08, 2010. This work was supported in part by the Engineering and Physical Sciences Research Council (EPSRC) of the U.K. under Grant GR/S27658/01, the Royal Society of the U.K., the National 973 Program of China under Grant 2009CB320600, and the Alexander von Humboldt Foundation of Germany. Recommended by Associate Editor G. Chesi.

Z. Wang is with the School of Information Science and Technology, Donghua University, Shanghai 200051, China. He is also with the Department of Information Systems and Computing, Brunel University, Uxbridge, Middlesex UB8 3PH, U.K. (e-mail: Zidong.Wang@brunel.ac.uk).

Y. Liu is with the Department of Mathematics, Yangzhou University, Yangzhou 225002, China.

X. Liu is with the Department of Information Systems and Computing, Brunel University, Uxbridge, Middlesex, UB8 3PH, U.K.

Color versions of one or more of the figures in this technical note are available online at <http://ieeexplore.ieee.org>.

Digital Object Identifier 10.1109/TAC.2010.2046114